

remain a frustrating unknown until considerably more research is done.

References

1. Stickgold, R., Whidbee, D., Schirmer, B., Patel, V., and Hobson, J.A. (2000). Visual discrimination task improvement: A multi-step process occurring during sleep. *J. Cogn. Neurosci.* 12, 246–254.
2. Frank, M.G., Issa, N.P., and Stryker, M.P. (2001). Sleep enhances plasticity in the developing visual cortex. *Neuron* 30, 275–287.
3. Jackson, C., McCabe, B.J., Nicol, A.U., Grout, A.S., Brown, M.W., and Horn, G. (2008). Dynamics of a memory trace: effects of sleep on consolidation. *Curr. Biol.* 18, 393–400.
4. Horn, G. (2004). Pathways of the past: the imprint of memory. *Nat. Rev. Neurosci.* 5, 108–120.
5. Horn, G., Nicol, A.U., and Brown, M.W. (2001). Tracking memory's trace. *Proc. Natl. Acad. Sci. USA* 98, 5282–5287.
6. Stickgold, R. (2005). Sleep-dependent memory consolidation. *Nature* 437, 1272–1278.
7. Ganguly-Fitzgerald, I., Donlea, J., and Shaw, P.J. (2006). Waking experience affects sleep need in *Drosophila*. *Science* 313, 1775–1781.
8. Smith, C. (1995). Sleep states and memory processes. *Behav. Brain Res.* 69, 137–145.
9. Nishida, M., and Walker, M.P. (2007). Daytime naps, motor memory consolidation and regionally specific sleep spindles. *PLoS ONE* 2, e341.
10. Huber, R., Ghilardi, M.F., Massimini, M., and Tononi, G. (2004). Local sleep and learning. *Nature* 430, 78–81.
11. Peigneux, P., Laureys, S., Fuchs, S., Collette, F., Perrin, F., Reggers, J., Phillips, C., Degueldre, C., Del Fiore, G., Aerts, J., *et al.* (2004). Are spatial memories strengthened in the human hippocampus during slow wave sleep? *Neuron* 44, 535–545.
12. Walker, M.P., Stickgold, R., Alsop, D., Gaab, N., and Schlaug, G. (2005). Sleep-dependent motor memory plasticity in the human brain. *Neurosci.* 133, 911–917.
13. Wilson, M.A., and McNaughton, B.L. (1994). Reactivation of hippocampal ensemble memories during sleep. *Science* 265, 676–679.
14. Dave, A.S., and Margoliash, D. (2000). Song replay during sleep and computational rules for sensorimotor vocal learning. *Science* 290, 812–816.
15. Manoach, D.S., Cain, M.S., Vangel, M.G., Khurana, A., Goff, D.C., and Stickgold, R. (2004). A failure of sleep-dependent procedural learning in chronic, medicated schizophrenia. *Biol. Psychiat.* 56, 951–956.
16. Bushey, D., Huber, R., Tononi, G., and Cirelli, C. (2007). *Drosophila* Hyperkinetic mutants have reduced sleep and impaired memory. *J. Neurosci.* 27, 5384–5393.
17. Vyazovskiy, V.V., Cirelli, C., Pfister-Genskow, M., Faraguna, U., and Tononi, G. (2008). Molecular and electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. *Nat. Neurosci.* 11, 200–208.
18. Pitman, J.L., McGill, J.J., Keegan, K.P., and Allada, R. (2006). A dynamic role for the mushroom bodies in promoting sleep in *Drosophila*. *Nature* 441, 753–756.

Department of Psychiatry, Harvard Medical School and Center for Sleep and Cognition, Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA.
E-mail: rstickgold@hms.harvard.edu

DOI: 10.1016/j.cub.2008.03.022

Sexual Dimorphism: Can You Smell the Difference?

A powerful new technique for visualizing neurons in the fly brain has uncovered fine neuroanatomical differences between the olfactory circuitries of male and female *Drosophila*.

Elizabeth J. Rideout
and Stephen F. Goodwin*

Dramatic behavioral differences exist between males and females in *Drosophila* [1–3]. Although the genetic basis for the creation of two morphologically and behaviorally distinct sexes has been extensively studied, few anatomical differences in the brain have been identified which may explain these dimorphic behaviors [4–7]. In fact, gross anatomical studies suggest that male and female brains are largely similar [5]. Identifying potential differences requires detailed analyses of the neural circuits underlying these sex-specific behaviors. Currently, identifying and tracing the projections of single neurons or populations of neurons relies upon clonal analysis techniques, which can be time-consuming due to the stochastic nature of generating uniquely labelled neurons [8,9].

Now, Datta *et al.* [10] have described a novel system for visualizing the cell bodies and projections of single neurons with high spatial and temporal resolution, using a photoactivatable

green fluorescent protein (PA-GFP) [11]. PA-GFP is a stable photoactivatable variant of GFP that after irradiation with 413 nm light shows a 100-fold increase in fluorescence when excited by 488 nm light [11]. Significantly, when PA-GFP is photoactivated in an isolated region of the cell, it will then diffuse throughout the entire cell. Datta *et al.* [10] exploited this property of PA-GFP to locate cell bodies of neurons, and to visualize axonal projections of identified neurons. Using this elegant technique, the authors were able to visualize and identify sexually dimorphic axonal projections of a specific population of neurons in the brain.

During courtship, male and female flies exchange a variety of stimulatory and inhibitory sensory cues [12]. Intriguingly, certain auditory and olfactory cues elicit opposite behavioral responses from the two sexes [13,14]; for example, *cis*-vaccenyl acetate, a male-specific pheromone, has been shown to have an inhibitory effect on male courtship, whereas in females, *cis*-vaccenyl acetate enhances receptivity to

copulation [14]. How can a single pheromone elicit such different behavioral responses? Datta *et al.* [10] addressed this question by investigating whether anatomical and/or functional differences in olfactory neural circuitry may explain this dimorphic behavior (Figure 1).

In flies, the physiological response to *cis*-vaccenyl acetate is mediated by a class of olfactory sensory neurons expressing the odorant receptor gene *Or67d*. All neurons expressing *Or67d* innervate a single glomerulus, DA1, in the antennal lobe [14]. Although the size of the DA1 glomerulus is sexually dimorphic [5,14,15], Datta *et al.* [10] found no essential differences between males and females in either the increase of intracellular Ca²⁺ concentration or in the electrophysiological responses in the DA1 glomerulus following exposure to *cis*-vaccenyl acetate, suggesting that the neurobiological basis for the sex-specific responses to *cis*-vaccenyl acetate must lie elsewhere in the olfactory neural circuit.

Next, using PA-GFP, they determined that projection neurons which project from the DA1 glomerulus to the lateral horn, a higher olfactory processing centre, have sexually dimorphic axonal arbors on the lateral horn (Figure 1). After identifying these sexually dimorphic arbors, Datta *et al.* [10] went on to show that this dimorphism depends on the expression of the male-specific isoform




Figure 1. Sexual dimorphism in higher olfactory processing centres.

The *Drosophila* olfactory circuitry. An odor is detected by an olfactory sensory neuron expressing a single type of odorant receptor gene (*Or*). All olfactory sensory neurons expressing a particular *Or* project their axons to a single glomerulus in the antennal lobe, where they form synapses with projection neurons, which transmit olfactory information to higher olfactory processing centres: the mushroom bodies and the lateral horn. Olfactory sensory neurons expressing *Or67d* largely govern the fly's physiological response to the male-specific pheromone *cis*-vaccenyl acetate [14], and project their axons to the DA1 glomerulus in the antennal lobe. DA1 is sexually dimorphic in volume, but no dimorphisms in the electrophysiological response to *cis*-vaccenyl acetate were detected in this glomerulus. Using photoactivatable green fluorescent protein (PA-GFP) to visualize both individual DA1 projection neurons, and subpopulations of DA1 projection neurons, Datta *et al.* [10] investigated whether a dimorphism occurs in the transmission of olfactory information to higher olfactory processing centres. A dimorphic axonal arbor of DA1 projection neurons on the lateral horn was uncovered (black arrowheads), where males, projection neurons had additional axonal branches. However, the impact of this dimorphism on behavior remains unknown (dashed line).

of *fruitless* (*fru*), a sex determination gene required for the performance of many male behaviors [1]. Previously, the dimorphism in DA1 glomerular volume had been shown to depend on *fru* [5]. These new findings extend our understanding of *fru*'s role in shaping sex-specific olfactory circuitry [10]. In the future, it will be interesting to see whether *fru*'s male-specific isoforms (*Fru^M*) are also responsible for the creation of other dimorphisms in olfactory circuitry, such as those recently reported by Jefferis *et al.* [16] in the lateral horn.

Although Datta *et al.* [10] were not able unequivocally to link these dimorphisms in DA1 projection neurons to specific behavioral outcomes, they have developed a strategy for targeted neural tracing that allows electrophysiological recordings to be taken from identified neurons. This approach will undoubtedly be exploited to identify additional dimorphic circuitry in

the *Drosophila* brain. It is important to mention, however, that in trying to catalogue sexual dimorphism in the brain, there are examples where equivalent neurons are not present in both sexes for comparison purposes [4–7]. In fact, one of these studies [4] showed that, for a cluster of neurons dorsal to the antennal lobe, both the presence of the neuron and its axonal morphology depend on the presence of *Fru^M* proteins. Together with the findings from Datta *et al.* [10], these results suggest that *fru*'s influence on male courtship behavior is a product of regulating dimorphism in the brain at many levels. Along with other sex determination genes, *fru* specifies dimorphic neuronal populations, and in addition, dimorphic axonal and dendritic projections, factors which all contribute to sex-specific behaviors.

So how does the identification of anatomical differences between males and females allow us to gain the

necessary insights into the origins of sexually dimorphic behaviors? First, the structure of the *Drosophila* brain has been well studied, and various regions of the brain are known to be associated with specific behaviors; for example, the ability to learn and remember is intimately linked with the mushroom bodies. Therefore, by understanding where the dimorphisms between males and females lie, the significance of any existing dimorphisms can be inferred, and later tested. Second, as Datta *et al.* [10] indicate, the ability to label specific neurons or specific subsets of neurons in the living brain will allow recordings to be taken from these identified neurons in flies (or indeed other animals) as they are exposed to specific cues, or even as they are behaving, allowing the function of a dimorphic neuron to be related to behavior.

Finally, experience-dependent changes in specific brain structures such as the mushroom bodies have been reported [17]; in the future, it will be interesting to see if this new system for visualizing specific structures and/or neurons can be used to investigate the dynamics of these experience-dependent changes in the brain, giving vital insights into the mechanisms underlying both synaptic and behavioral plasticity. Therefore, this elegant new technique for visualizing and recording from specific neurons moves us another step closer to understanding the relationship between the development and function of the brain, and behavior.

References

1. Billeter, J.C., Rideout, E.J., Dorman, A.J., and Goodwin, S.F. (2006). Control of male sexual behavior in *Drosophila* by the sex determination pathway. *Curr. Biol.* 16, R766–R776.
2. Nilsen, S.P., Chan, Y.B., Huber, R., and Kravitz, E.A. (2004). Gender-selective patterns of aggressive behavior in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* 101, 12342–12347.
3. Martin, J.R., Ernst, R., and Heisenberg, M. (1999). Temporal pattern of locomotor activity in *Drosophila melanogaster*. *J. Comp. Physiol. A* 184, 73–84.
4. Kimura, K., Ote, M., Tazawa, T., and Yamamoto, D. (2005). *Fruitless* specifies sexually dimorphic neural circuitry in the *Drosophila* brain. *Nature* 438, 229–233.
5. Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirian, L., and Dickson, B.J. (2005). Neural circuitry that governs *Drosophila* male courtship behavior. *Cell* 121, 795–807.
6. Billeter, J.C., Villella, A., Allendorfer, J.B., Dorman, A.J., Richardson, M., Gailey, D.A., and Goodwin, S.F. (2006). Isoform-specific control of male neuronal differentiation and behavior in *Drosophila* by the *fruitless* gene. *Curr. Biol.* 16, 1063–1076.

- Rideout, E.J., Billeter, J.C., and Goodwin, S.F. (2007). The sex-determination genes *fruitless* and *doublesex* specify a neural substrate required for courtship song. *Curr. Biol.* 17, 1473–1478.
- Lee, T., and Luo, L. (2001). Mosaic analysis with a repressible cell marker (MARCM) for *Drosophila* neural development. *Trends Neurosci.* 24, 251–254.
- Wong, A.M., Wang, J.W., and Axel, R. (2002). Spatial representation of the glomerular map in the *Drosophila* protocerebrum. *Cell* 109, 229–241.
- Datta, S.R., Vasconcelos, M.L., Ruta, V., Luo, S., Wong, A., Demir, E., Flores, J., Balonze, K., Dickson, B.J., and Axel, R. (2008). The *Drosophila* pheromone cVA activates a sexually dimorphic neural circuit. *Nature* 452, 473–477.
- Patterson, G.H., and Lippincott-Schwartz, J. (2002). A photoactivatable GFP for selective photolabelling of proteins and cells. *Science* 297, 1873–1877.
- Ferveur, J.F. (2005). Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal communication. *Behav. Genet.* 35, 279–295.
- von Schilcher, F. (1976). The role of auditory stimuli in the courtship of *Drosophila melanogaster*. *Anim. Behav.* 24, 18–26.
- Kurtovic, A., Widmer, A., and Dickson, B.J. (2007). A single class of olfactory neurons mediates behavioural responses to a *Drosophila* sex pheromone. *Nature* 446, 542–546.
- Kondoh, Y., Kaneshiro, K.Y., Kimura, K., and Yamamoto, D. (2003). Evolution of sexual dimorphism in the olfactory brain of Hawaiian *Drosophila*. *Proc. Biol. Sci.* 270, 1005–1013.
- Jefferis, G.S., Potter, C.J., Chan, A.M., Marin, E.C., Rohlffing, T., Maurer, C.R. Jr., and Luo, L. (2007). Comprehensive maps of *Drosophila* higher olfactory centers: spatially segregated fruit and pheromone representation. *Cell* 128, 1187–1203.
- Technau, G.M. (1984). Fiber number in the mushroom bodies of adult *Drosophila melanogaster* depends on age, sex and experience. *J. Neurogenet.* 1, 113–126.

University of Glasgow, IBLs-Division of Molecular Genetics, Anderson College, 56 Dumbarton Road, Glasgow G11 6NU, UK.
*E-mail: s.goodwin@bio.gla.ac.uk

DOI: 10.1016/j.cub.2008.03.035

Visual Categorization: When Categories Fall to Pieces

We cannot help but categorize the visual world into objects like cats and faces. An intriguing new study shows that observers automatically discover informative fragments of visual objects during category learning.

Quoc C. Vuong

We see the world in discrete categories in order to recognize and interact appropriately with objects in our environment [1]. How do we learn visual object categories? Our intuition suggests that, through experience, we acquire features found in members of one category but not in those from another category. For example, cats have whiskers; human faces, on the other hand, normally do not. There is empirical support for this intuitive view [2,3].

But a fundamental problem with this intuition is image variability. Familiar objects from the same category can have an enormous range of appearance; they are often occluded by other objects; how they appear to us can further be confounded by viewing conditions such as variable illumination; and so on [2]. These factors converge to make it extremely difficult to learn generic features that are reliable for visual categorization.

In work published recently in *Current Biology*, Hegd  et al. [4] offer a compelling solution to this problem, but one that highlights the need for us to re-think the pieces that make up objects and object categories. Armed with a set of novel visual categories [5] and a statistical means to select features [6,7], these authors have demonstrated that

observers automatically discover fragments — literally, bits and pieces of images — during category learning that are very effective for visual categorization. This provides a new and important link between visual category learning and visual categorization.

In this new study [4], observers classified a large number of unfamiliar objects into two categories. The

objects were synthesized from a novel virtual phylogenesis algorithm which simulated the evolution of biological forms [5], so that category members captured natural variations of categories we are more familiar with. The examples in Figure 1 show that this classification task is far from trivial, even with whole objects (see supplemental Figure S1 in the paper for more examples).

Two main sets of image fragments were extracted from trained objects using the same statistical procedure. Observers then classified all fragments, just as they had done with whole objects. This sounds like an even more daunting task. Amazingly though, observers were as accurate with one set of fragments as they were




Figure 1. Example objects synthesized by virtual phylogenesis. Observers were only trained on objects from two of the three categories A, B and C (from [4]).